

Generating solutions of "impossible-to-solve" problems and simulating "impossible-to-simulate" models

Florent Krzakala
Espci-Paristech

L. Zdeborová (Los Alamos, LANL)

Planting !


Florent Krzakala
Espci-Paristech

L. Zdeborová (Los Alamos, LANL)

Impossible-to-solve problems

Some optimization problems such as COL and SAT are almost impossible to solve!

ex: Hard Instances of random graph coloring




A 3-coloring of a random graph with $c=4.36$

Impossible-to-solve problems

Some optimization problems such as COL and SAT are almost impossible to solve!

ex: Hard Instances of random graph coloring




A 3-coloring of a random graph with $c=4.36$

The “ $q \log q$ ” problem

Impossible-to-solve problems

Some optimization problems such as COL and SAT are almost impossible to solve!

ex: Hard Instances of random graph coloring



A 3-coloring of a random graph with $c=4.36$

The “ $q \log q$ ” problem

- Consider q color (with q large enough) and a large random graph of average degree c
- W.h.p this graph is colorable if $c < 2q \log q$
- However, no algorithm is able to do so efficiently (polynomial) for $c > q \log q$!

D. Achlioptas et al. Nature 2005

$q \log q$


$2q \log q$

Average
degree c

Impossible-to-solve problems

Some optimization problems such as COL and SAT are almost impossible to solve!

ex: Hard Instances of random graph coloring




A 3-coloring of a random graph with $c=4.36$

The “ $q \log q$ ” problem

- Consider q color (with q large enough) and a large random graph of average degree c
- W.h.p this graph is colorable if $c < 2q \log q$
- However, no algorithm is able to do so efficiently (polynomial) for $c > q \log q$!


D. Achlioptas et al. Nature 2005



Impossible-to-solve problems

Some optimization problems such as COL and SAT are almost impossible to solve!

ex: Hard Instances of random graph coloring




A 3-coloring of a random graph with $c=4.36$

The “ $q \log q$ ” problem

- Consider q color (with q large enough) and a large random graph of average degree c
- W.h.p this graph is colorable if $c < 2q \log q$
- However, no algorithm is able to do so efficiently (polynomial) for $c > q \log q$!


D. Achlioptas et al. Nature 2005



Impossible-to-solve problems

Some optimization problems such as COL and SAT are almost impossible to solve!

ex: Hard Instances of random graph coloring



A 3-coloring of a random graph with $c=4.36$


The “ $q \log q$ ” problem

- Consider q color (with q large enough) and a large random graph of average degree c
- W.h.p this graph is colorable if $c < 2q \log q$
- However, no algorithm is able to do so efficiently (polynomial) for $c > q \log q$!



No one has ever seen the solution of, say 5-coloring, for large enough c and $N=10^6$

D. Achlioptas et al. Nature 2005



Impossible-to-simulate problems

Some optimization problems such as COL and SAT are also hard to sample

$q \log q$

$2q \log q$

*Average
degree c*

Impossible-to-simulate problems

Some optimization problems such as COL and SAT are also hard to sample

*Consider the following “coloring”
or “Potts-Antiferromagnet”
Hamiltonian*

$$\mathcal{H} = \sum_{\langle ij \rangle} \delta(s_i, s_j)$$
$$s_i = 1, 2, \dots, q$$

$q \log q$

$2q \log q$

*Average
degree c*

Impossible-to-simulate problems

Some optimization problems such as COL and SAT are also hard to sample

*Consider the following “coloring”
or “Potts-Antiferromagnet”
Hamiltonian*

$$\mathcal{H} = \sum_{\langle ij \rangle} \delta(s_i, s_j)$$
$$s_i = 1, 2, \dots, q$$


Temperature

Dynamic transition

$q \log q$

$2q \log q$

*Average
degree c*




Impossible-to-simulate problems

Some optimization problems such as COL and SAT are also hard to sample

Consider the following “coloring”
or “Potts-Antiferromagnet”
Hamiltonian


$$\mathcal{H} = \sum_{\langle ij \rangle} \delta(s_i, s_j)$$
$$s_i = 1, 2, \dots, q$$




Frustrating Intractable Problems

- * We know that some random problems DO have solutions, but we cannot find them!
- * Sampling and performing Monte-Carlo is even Harder!
- * Many predictions from statistical physics in random problems.... but impossible to test most of them !


A Different “Reverse” strategy



A Different “Reverse” strategy




A Different “Reverse” strategy



Instead of choosing a problem, and looking for a solution....

A Different “Reverse” strategy



Instead of choosing a problem, and looking for a solution....




*We choose a configuration/assignment and
and look for a problem for which this is a solution !*

The Planted Ensemble in the coloring problem

Consider the 3-coloring problem with N nodes and M links.


- 1) Color randomly
the N nodes



The Planted Ensemble in the coloring problem

Consider the 3-coloring problem with N nodes and M links.


- i) Color randomly the N nodes
- ii) Put the M links randomly such that the planted configuration is a proper coloring



The Planted Ensemble in the coloring problem

Consider the 3-coloring problem with N nodes and M links.

- I) Color randomly the N nodes
- II) Put the M links randomly such that the planted configuration is a proper coloring
- III) Now, we have created a problem for which **we know** the solution




The Planted Ensemble in the coloring problem

Consider the 3-coloring problem with N nodes and M links.

I) Color randomly
the N nodes

II) Put the M links randomly such
that the planted configuration is a
proper coloring

III) Now, we have created a problem
for which **we know** the solution



IV) We could also have prepared a configuration with a known cost/energy

The Random ensemble versus the Planted ensemble

Random ensemble

Choose a random graph
with N nodes and M links

Planted ensemble

Choose a random coloring of N
nodes

Choose a random graph such
that this is a correct coloring...

The Random ensemble versus the Planted ensemble

Random ensemble

Choose a random graph with N nodes and M links

Planted ensemble

Choose a random coloring of N nodes

Choose a random graph such that this is a correct coloring...

- * Is it really the same to look for a solution a random problem and to look for a random problem that matches a random solution ?

The Random ensemble versus the Planted ensemble

Random ensemble

Choose a random graph with N nodes and M links

Planted ensemble

Choose a random coloring of N nodes

Choose a random graph such that this is a correct coloring...

- * Is it really the same to look for a solution a random problem and to look for a random problem that matches a random solution ?
- * The surprising answer is: in some cases **YES!**

The Random ensemble versus the Planted ensemble

Random ensemble

Choose a random graph with N nodes and M links

Planted ensemble

Choose a random coloring of N nodes

Choose a random graph such that this is a correct coloring...

- * Is it really the same to look for a solution a random problem and to look for a random problem that matches a random solution ?
- * The surprising answer is: in some cases **YES!**



Montanari and Semerjian, Jstat. '06 & Achlioptas and Coja-Oghlan, arXiv:0803.2122:

The two ensembles are asymptotically ($N \rightarrow \infty$) equivalent for low enough degree c !

The Random ensemble versus the Planted ensemble

Random ensemble

Choose a random graph with N nodes and M links

Planted ensemble

Choose a random coloring of N nodes

Choose a random graph such that this is a correct coloring...

Definition : Two ensembles of random graphs are asymptotically equivalent if and only if in the thermodynamic limit every property which is almost surely true on a graph from one ensemble is also almost surely true on a graph from the other ensemble.

Some open questions:

- * Until which connectivity/degree c the planted and random ensembles are equivalent ?
- * Is the planted ensemble interesting beyond this connectivity ?
- * Can we generalize this approach to finite energy (coloring with a finite fraction of mistakes ?)
- * How can we use a planted configuration ?
- * What are the models where a “quiet” planting is possible ?

In this talk:

- 1) A (brief) summary of a theory of “quiet” planting in random models
- 2) Using planted configurations for fast simulations.

The Planted Ensemble

The Planted Ensemble

* We use the formalism described in Zdeborová's talk

Main result

Consider a model where the annealed computation is correct in some region (high temperature or low degree)

$$f = -\frac{1}{N}\beta[\log Z]_{dis} \longleftrightarrow f_{annealed} = -\frac{1}{N}\beta \log [Z]_{dis}$$

Main result

Consider a model where the annealed computation is correct in some region (high temperature or low degree)

$$f = -\frac{1}{N}\beta[\log Z]_{dis} \longleftrightarrow f_{annealed} = -\frac{1}{N}\beta \log [Z]_{dis}$$

=

Consider a model where a factorized (i.e. identical for all nodes)
Belief Propagation solution is correct in some region
(high temperature or low degree)

A list of models with “Quiet” planting !

This condition is fulfilled (at least in some region) for many models:

- Random q -coloring problem
- Random XOR-SAT
- Mean field spin glasses (e.g. Vianna-Bray, Sherrington-Kirkpatrick)
- Random 2-in-4 Sat
- Random Vertex-Cover (independent set)
- Any non disordered model on a random regular graphs
-

A list of models with “Quiet” planting !

This condition is fulfilled (at least in some region) for many models:

- Random q-coloring problem
- Random XOR-SAT
- Mean field spin glasses (e.g. Vianna-Bray, Sherrington-Kirkpatrick)
- Random 2-in-4 Sat
- Random Vertex-Cover (independent set)
- Any non disordered model on a random regular graphs
-

This condition is not fulfilled for :

- Random K-SAT

Main result

Consider a model where the annealed computation is correct in some region (high temperature or low degree)

=

Consider a model where a factorized (i.e. identical for all nodes) Belief Propagation solution is correct in some region (high temperature or low degree)

Main result

Consider a model where the annealed computation is correct in some region (high temperature or low degree)

=

Consider a model where a factorized (i.e. identical for all nodes) Belief Propagation solution is correct in some region (high temperature or low degree)



**In the region where the
two free energies are equal,
the two ensembles are equivalent**

Main result

Consider a model where the annealed computation is correct in some region (high temperature or low degree)



Consider a model where a factorized (i.e. identical for all nodes) Belief Propagation solution is correct in some region (high temperature or low degree)



In the region where the two free energies are equal, the two ensembles are equivalent



In the region where the two free energies are different, the planted configuration induces an additional “Gibbs” state (or BP fixed point)

A list of models with “Quiet” planting !

This condition is fulfilled (at least in some region) for many models:

- Random q -coloring problem
- Random XOR-SAT
- Mean field spin glasses (e.g. Vianna-Bray, Sherrington-Kirkpatrick)
- Random 2-in-4 Sat
- Random Vertex-Cover (independent set)
- Any non disordered model on a random regular graphs
-

A list of models with “Quiet” planting !

This condition is fulfilled (at least in some region) for many models:

- Random q-coloring problem
- Random XOR-SAT
- Mean field spin glasses (e.g. Vianna-Bray, Sherrington-Kirkpatrick)
- Random 2-in-4 Sat
- Random Vertex-Cover (independent set)
- Any non disordered model on a random regular graphs
-

For all these models, the cavity method allows to compute the value of the threshold beyond which $f \neq f_{\text{annealed}}$

⇒ “Phase transition”

A list of models with “Quiet” planting !

This condition is fulfilled (at least in some region) for many models:

- Random q-coloring problem
- Random XOR-SAT
- Mean field spin glasses (e.g. Vianna-Bray, Sherrington-Kirkpatrick)
- Random 2-in-4 Sat
- Random Vertex-Cover (independ
- Any non disordered model on a random regular graphs
-

*In some models, the
equivalence can be proven*

For all these models, the cavity method allows to compute
the value of the threshold beyond which $f \neq f_{\text{annealed}}$


⇒ “Phase transition”

Take-home message:

- * Conjecture 1: the planted model is equivalent to the original one up to the point where the annealed solution is correct
(for physicists: up to the static spin glass transition...) and the planted configuration is a “typical” one.


A Solution To an Impossible-To-Solve Problem

ex: 5-coloring of Erdős-Renyi random graphs



A Solution To an Impossible-To-Solve Problem

ex: 5-coloring of Erdős-Renyi random graphs




A Solution To an Impossible-To-Solve Problem


ex: 5-coloring of Erdős-Renyi random graphs

5-coloring using walkcol with

$$N=10^6$$




*One can create impossible to solve
problems of any size
where the solution is known
only by the creator*



A Solution To an Impossible-To-Solve Problem


ex: q -coloring of Erdős-Renyi random graphs for large q



A Solution To an Impossible-To-Solve Problem

ex: q -coloring of Erdős-Renyi random graphs for large q

Planting !




Take-home message:

- * Conjecture 1: the planted model is equivalent to the original one up to the point where the annealed solution is correct
(for physicists: up to the spin glass transition...) and the planted configuration is a “typical” one

Take-home message:

- * Conjecture 1: the planted model is equivalent to the original one up to the point where the annealed solution is correct
(for physicists: up to the spin glass transition...) and the planted configuration is a “typical” one

*U. Feige, E. Mossel and D. Vilenchik.
Proceedings of Random'06, LNCS 4410,*



*Planted configuration easy
to find for large enough c*

Take-home message:


- * Conjecture 1: the planted model is equivalent to the original one up to the point where the annealed solution is correct

(for physicists: up to the spin glass transition...)
and the planted configuration is a “typical” one

- * Conjecture 2: Planted configuration are hard to find until the so-called Kesten-Stigum threshold,
(for physicists: this is the local spin glass instability)
beyond which they can be solved easily using BP.


Example in coloring

The planted solution is well hidden until $c_{KS} = (q-1)^2$



Example in coloring

The planted solution is well hidden until $c_{KS} = (q-1)^2$




Example in coloring

The planted solution is well hidden until $c_{KS} = (q-1)^2$



Example in coloring

The planted solution is well hidden until $c_{KS} = (q-1)^2$




Simulating “impossible-to-simulate” models

*How to perform simulations that
are usually considered to be impossible?*

Impossible-to-simulate problems

Random optimization problems & mean-field spin glasses


Temperature



*Average
degree c*


Impossible-to-simulate problems

Random optimization problems & mean-field spin glasses



Impossible-to-simulate problems

Random optimization problems & mean-field spin glasses




Modus operandi

1. Plant a configuration, create the graph such that the configuration satisfies all constraints
2. We now have a random instance and a “typical” equilibrium solution at zero temperature
3. We use it !

Example 1:

Testing the cavity predictions for the clustering transition



$$\psi_{factorized} = \left(\frac{1}{q}, \frac{1}{q} \cdots \frac{1}{q} \right)$$


$$M = \frac{1}{qN} \sum_{graph} \sum_{c=1}^q \frac{\psi_{BP}^{c,i} - \frac{1}{q}}{1 - \frac{1}{q}}$$

FK, Montanari, Semerjian, Ricci-Tersenghi, Zdeborova, PNAS 07 & FK and Zdeborova, PRE 07

Prediction: beyond the so-called “dynamic” threshold, a non-trivial non-factorized fixed point of BP is obtained if one starts from an equilibrium configuration

Example 1:

Testing the cavity predictions for the clustering transition



average degree

$$\psi_{factorized} = \left(\frac{1}{q}, \frac{1}{q} \dots \frac{1}{q} \right)$$

$$M = \frac{1}{qN} \sum_{graph} \sum_{c=1}^q \frac{\psi_{BP}^{c,i} - \frac{1}{q}}{1 - \frac{1}{q}}$$

Simulation with $N=10^6$


Prediction: beyond the so-called “dynamic” threshold, a non-trivial non-factorized fixed point of BP is obtained if one starts from an equilibrium configuration

Modus operandi for finite temperature simulations

1. Plant a configuration, create the graph such that the configuration has exactly the equilibrium energy
2. We now have a random instance and a “typical” equilibrium solution at temperature T
3. We use it !

Example 2:

Testing the cavity predictions for the clustering transition



Prediction: beyond the so-called “dynamic” threshold, the Monte-Carlo Dynamic is trapped!

Ex: 3-XORSAT, $T=0.255, c=3$

Example 2:

Testing the cavity predictions for the clustering transition

Usual Approach:

- 1) *Start with a random initial condition*
- 2) *compute the correlation function:*


$$C(t) = \frac{1}{N} \sum_{i=1}^N S_i(t_{init} = 0) S_i(t)$$

Prediction: beyond the so-called “dynamic” threshold, the Monte-Carlo Dynamic is trapped!

Ex: 3-XORSAT, $T_d=0.255$

Example 2:

Testing the cavity predictions for the clustering transition



Usual Approach:

- 1) Start with a random initial condition
- 2) compute the correlation function:


$$C(t) = \frac{1}{N} \sum_{i=1}^N S_i(t_{init} = 0) S_i(t)$$

Prediction: beyond the so-called “dynamic” threshold, the Monte-Carlo Dynamic is trapped!

Ex: 3-XORSAT, $T_d=0.255$

Example 2:

Testing the cavity predictions for the clustering transition



Usual Approach:

- 1) Start with a random initial condition
- 2) Try to find an equilibrium configuration
- 2) compute the correlation function:


$$C(t) = \frac{1}{N} \sum_{i=1}^N S_i(t_{init} = t_w) S_i(t - t_w)$$

Prediction: beyond the so-called “dynamic” threshold, the Monte-Carlo Dynamic is trapped!

Ex: 3-XORSAT, $T_d=0.255$

Example 2:

Testing the cavity predictions for the clustering transition



Usual Approach:

- 1) Start with a random initial condition
- 2) Try to find an equilibrium configuration
- 2) compute the correlation function:

$$C(t) = \frac{1}{N} \sum_{i=1}^N S_i(t_{init} = t_w) S_i(t - t_w)$$


tw=10

Prediction: beyond the so-called “dynamic” threshold, the Monte-Carlo Dynamic is trapped!

Ex: 3-XORSAT, $T_d=0.255$

Example 2:

Testing the cavity predictions for the clustering transition



Usual Approach:

- 1) Start with a random initial condition
- 2) Try to find an equilibrium configuration
- 2) compute the correlation function:

$$C(t) = \frac{1}{N} \sum_{i=1}^N S_i(t_{init} = t_w) S_i(t - t_w)$$


tw=100

Prediction: beyond the so-called “dynamic” threshold, the Monte-Carlo Dynamic is trapped!

Ex: 3-XORSAT, $T_d=0.255$

Example 2:

Testing the cavity predictions for the clustering transition



Usual Approach:

- 1) Start with a random initial condition
- 2) Try to find an equilibrium configuration
- 2) compute the correlation function:

$$C(t) = \frac{1}{N} \sum_{i=1}^N S_i(t_{init} = t_w) S_i(t - t_w)$$


tw=1000

Prediction: beyond the so-called “dynamic” threshold, the Monte-Carlo Dynamic is trapped!

Ex: 3-XORSAT, $T_d=0.255$

Example 2:

Testing the cavity predictions for the clustering transition



Usual Approach:

- 1) Start with a random initial condition
- 2) Try to find an equilibrium configuration
- 2) compute the correlation function:

$$C(t) = \frac{1}{N} \sum_{i=1}^N S_i(t_{init} = t_w) S_i(t - t_w)$$


tw=10000

Prediction: beyond the so-called “dynamic” threshold, the Monte-Carlo Dynamic is trapped!

Ex: 3-XORSAT, $T_d=0.255$

Example 2:

Testing the cavity predictions for the clustering transition



A better Approach:

- 1) Start with an equilibrated initial condition
- 2) compute the correlation function:


$$C(t) = \frac{1}{N} \sum_{i=1}^N S_i(t_{init} = 0) S_i(t)$$

Prediction: beyond the so-called “dynamic” threshold, the Monte-Carlo Dynamic is trapped!

Ex: 3-XORSAT, $T_d=0.255$

Example 2:

Testing the cavity predictions for the clustering transition



A better Approach:

- 1) Start with an equilibrated initial condition
- 2) compute the correlation function:


$$C(t) = \frac{1}{N} \sum_{i=1}^N S_i(t_{init} = 0) S_i(t)$$

Prediction: beyond the so-called “dynamic” threshold, the Monte-Carlo Dynamic is trapped!

Ex: 3-XORSAT, $T_d=0.255$

Example 2:

Testing the cavity predictions for the clustering transition



A better Approach:

Start with an equilibrated initial condition
Many temperatures:

*Divergence of the
relaxation time*

Prediction: beyond the so-called “dynamic” threshold, the Monte-Carlo Dynamic is trapped!


Ex: 3-XORSAT, $T_d=0.255$

Example 3:

Studying Monte-Carlo annealings starting from equilibrium

XOR-SAT problems
(Parity-check)

$$\mathcal{H}(\{S\}) = \sum_{ijk} \frac{1 + J_{ijk} S_i S_j S_k}{2}$$




Prediction: cf. Zdeborová's talk:

Monte Carlo cooling and heating follow a well defined line

Example 3:

Studying Monte-Carlo annealings starting from equilibrium



XOR-SAT problems
(Parity-check)

$$\mathcal{H}(\{S\}) = \sum_{ijk} \frac{1 + J_{ijk} S_i S_j S_k}{2}$$


N=200 000 spins

Prediction: cf. Zdeborová's talk:

Monte Carlo cooling and heating follow a well defined line

Example 3:

Studying Monte-Carlo annealings starting from equilibrium



XOR-SAT problems
(Parity-check)

$$\mathcal{H}(\{S\}) = \sum_{ijk} \frac{1 + J_{ijk} S_i S_j S_k}{2}$$


N=200 000 spins

Prediction: cf. Zdeborová's talk:

Monte Carlo cooling and heating follow a well defined line

Example 3:

Studying Monte-Carlo annealings starting from equilibrium



XOR-SAT problems
(Parity-check)

$$\mathcal{H}(\{S\}) = \sum_{ijk} \frac{1 + J_{ijk} S_i S_j S_k}{2}$$


N=200 000 spins

Prediction: cf. Zdeborová's talk:

Monte Carlo cooling and heating follow a well defined line

Example 3:

Studying Monte-Carlo annealings starting from equilibrium



XOR-SAT problems
(Parity-check)

$$\mathcal{H}(\{S\}) = \sum_{ijk} \frac{1 + J_{ijk} S_i S_j S_k}{2}$$

N=200 000 spins

Prediction: cf. Zdeborová's talk:

Monte Carlo cooling and heating follow a well defined line

Example 4:

Studying more complex Hamiltonians at low temperature

$$\mathcal{H}(\{S\}) = \sum_{ijk} \frac{1 + J_{ijk} S_i S_j S_k}{2}$$

Example 4:

Studying more complex Hamiltonians at low temperature

$$\mathcal{H}(\{S\}) = \sum_{ijk} \frac{1 + J_{ijk} S_i S_j S_k}{2} + \Gamma \mathcal{H}_{\text{perturb}}$$

*Start with an equilibrated
configuration at $\Gamma=0$ and increase Γ*

Example 4:

Studying more complex Hamiltonians at low temperature


Example: include a quantum transverse field!

$$\mathcal{H}(\{S\}) = \sum_{ijk} \frac{1 + J_{ijk} S_i S_j S_k}{2} \quad \Rightarrow \quad \mathcal{H} = \sum_{ijk} \frac{1 + J_{ijk} s_i^z s_j^z s_k^z}{2} + \Gamma \sum_i s_i^x$$

Example 4:

Studying more complex Hamiltonians at low temperature


$$\mathcal{H} = \sum_{ijk} \frac{1 + J_{ijk} s_i^z s_j^z s_k^z}{2} + \Gamma \sum_i s_i^x$$



Example 4:

Studying more complex Hamiltonians at low temperature

$$\mathcal{H} = \sum_{ijk} \frac{1 + J_{ijk} s_i^z s_j^z s_k^z}{2} + \Gamma \sum_i s_i^x$$



***First order
Quantum
transition***



Imply the failure of
Quantum Annealing
(or Quantum Adiabatic
Algorithm)

Conclusions

A quiet planting is possible in many models.

- “Quiet” Planting does not change the properties of the ensemble up to the condensation threshold.
- Planted solutions are hard to find until the Kesten-Stigum threshold.
- Possibility to hide solutions (even a unique solution)

FK and L. Zdeborová:

* *Phys. Rev. Lett. 102, 238701 (2009)*

* *arXiv:0902.4185, submitted in SIAM Journal on Discrete Mathematics*

* *And more to come...*

Conclusions

A quiet planting is possible in many models.

- “Quiet” Planting does not change the properties of the ensemble up to the condensation threshold.
- Planted solutions are hard to find until the Kesten-Stigum threshold.
- Possibility to hide solutions (even a unique solution)

There **is** a free lunch: instantaneous simulations.

- Many “mean field” models and random optimization models can be simulated efficiently using planting at zero or finite temperature.

FK and L. Zdeborová:

** Phys. Rev. Lett. 102, 238701 (2009)*

** arXiv:0902.4185, submitted in SIAM Journal on Discrete Mathematics*

** And more to come...*